

Physics 6433, Quantum Field Theory
Assignment #6
Due Friday, October 23, 2009

October 16, 2009

1. Derive the following form for the Minkowski-space propagation function in the vicinity of the light-cone

$$\begin{aligned} m^2|(x - x')^2| \ll 1 : \quad \Delta_+(x - x') &\approx \frac{i}{4\pi^2} \frac{1}{(x - x')^2 + i\epsilon} \\ &= \frac{i}{4\pi^2} P \frac{1}{(x - x')^2} + \frac{1}{4\pi} \delta((x - x')^2), \end{aligned}$$

where P stands for the Cauchy principal value.

2. Derive the following asymptotic forms for the Minkowski-space propagation function for large spacelike, and large timelike intervals, respectively,

$$\begin{aligned} R = [(x - x')^2]^{1/2}, mR \gg 1 : \quad \Delta_+(x - x') &\approx i \frac{(2m)^{1/2}}{(4\pi R)^{3/2}} e^{-mR}, \\ T = [-(x - x')^2]^{1/2}, mT \gg 1 : \quad \Delta_+(x - x') &\approx e^{-i\pi/4} \frac{(2m)^{1/2}}{(4\pi T)^{3/2}} e^{-imT}. \end{aligned}$$

Discuss the correspondence between the Euclidean and Minkowskian propagation functions in these limits.