

# Physics 2414, Spring 2008

## Group Exercise 9, Apr 3, 2008

Name 1: \_\_\_\_\_ OUID 1: \_\_\_\_\_  
Name 2: \_\_\_\_\_ OUID 2: \_\_\_\_\_  
Name 3: \_\_\_\_\_ OUID 3: \_\_\_\_\_  
Name 4: \_\_\_\_\_ OUID 4: \_\_\_\_\_

Section Number: \_\_\_\_

### Conservation of Linear Momentum

**Conservation of Energy:** The sum of change in kinetic energy  $\Delta K$  and the change in potential energy  $\Delta U$  equals the work done by the non-conservative forces.

$$\Delta K + \Delta U = \Sigma W_{nc}. \quad (1)$$


**Momentum:** A mass  $m$  moving with a velocity  $\vec{v}$  has a momentum  $\vec{p} = m\vec{v}$  associated with it.

**Conservation of linear momentum:** If there is no external forces acting on a system, the total linear momentum of the system is conserved.

$$\text{If } \vec{F}_{\text{net}} = 0, \quad \vec{p}_{\text{tot}} = \vec{p}_{\text{itot}}. \quad (2)$$

### Problems

A skier with mass  $M_1 = 75$  kg starts from rest at point '1' at a height  $H = 50$  meters from the ground. Assume frictionless surfaces all throughout this problem.




1. *Before collision:*

Find the velocity  $v_2$  of the skier when he reaches point '2'.

2. *Conservation of linear momentum:*

The skier keeps moving on the horizontal frictionless surface with velocity  $v_2$ . The skier collides into a box of mass  $m_2 = 25$  kg at point 'c'. After the collision the skier grabs the box and then moves with the box as a single entity with velocity  $v_3$ .



(a) What is the initial momentum  $\vec{p}_{1i}$  of the skier just before the collision?

$$\vec{p}_{1i} = \quad (3)$$

(b) What is the initial momentum  $\vec{p}_{2i}$  of the box just before the

collision?

$$\vec{\mathbf{p}}_{2i} = \quad (4)$$

(c) What is the final momentum  $\vec{\mathbf{p}}_{1f}$  of the skier just after the collision in terms of  $v_3$ ?

$$\vec{\mathbf{p}}_{1f} = \quad (5)$$

(d) What is the final momentum  $\vec{\mathbf{p}}_{2f}$  of the box just after the collision in terms of  $v_3$ ?

$$\vec{\mathbf{p}}_{2f} = \quad (6)$$

(e) Is there any net external force acting on the system consisting of (skier+box)? (Is the system accelerating? Remember Newton's law,  $\vec{\mathbf{F}}_{\text{net}} = m\vec{\mathbf{a}}$ .)

$$\vec{\mathbf{F}}_{\text{net}} = \quad (7)$$

(f) During the collision the masses exert forces on each other. The skier exerts a force  $\vec{\mathbf{F}}_{21}$  on the box and the box exerts a force  $\vec{\mathbf{F}}_{12}$  on the skier. What can you conclude about the magnitude of the forces  $|\vec{\mathbf{F}}_{21}|$  and  $|\vec{\mathbf{F}}_{12}|$ ?

$$(i) |\vec{\mathbf{F}}_{21}| > |\vec{\mathbf{F}}_{12}|$$

$$(ii) |\vec{\mathbf{F}}_{21}| = |\vec{\mathbf{F}}_{12}|$$

$$(iii) |\vec{\mathbf{F}}_{21}| < |\vec{\mathbf{F}}_{12}|$$

Thus conclude that the internal forces cancel each other.

(g) Using (a) to (f) determine the total initial momentum  $\vec{\mathbf{p}}_{\text{tot}}$  and the total final momentum  $\vec{\mathbf{p}}_{f\text{tot}}$ .

$$\vec{\mathbf{p}}_{\text{tot}} = \vec{\mathbf{p}}_{1i} + \vec{\mathbf{p}}_{2i} = \quad (8)$$

$$\vec{\mathbf{p}}_{f\text{tot}} = \vec{\mathbf{p}}_{1f} + \vec{\mathbf{p}}_{2f} = \quad (9)$$


(h) Conservation of linear momentum states that

$$\text{if } \vec{\mathbf{F}}_{\text{net}} = 0, \quad \vec{\mathbf{p}}_{f\text{tot}} = \vec{\mathbf{p}}_{\text{tot}}. \quad (10)$$

Use this to determine the velocity  $v_3$  of (skier+box) just after the collision.

3. *After collision:*

The skier and the box keep moving up the hill as a single entity. They rise to a maximum height  $h$  above the horizontal surface.



(a) Find the height  $h$ .

4. *Analysis:*

(a) After the collision the skier and the box are moving as a single entity with velocity  $v_3$  at point '3'. Pick the correct answer.

- (i) Momentum of the skier is greater than the momentum of the box at point '3'.
- (ii) Momentum of the skier is less than the momentum of the box at point '3'.

(iii) Momentum of the skier is equal to the momentum of the box at point '3'.

(b) Is the kinetic energy conserved between point '2' and point '3'?

(i) Yes

(ii) No

Explain your answer.