

## [ Atomic, Molecular, Optical, and Chemical Physics ]

The Atomic, Molecular, Optical and Chemical Physics group studies the interactions between and the manipulation of atoms, molecules, electrons and photons at low temperatures and low energies. Current specific research areas in theory and experiment include chemical reaction dynamics, the physics of ultra-cold atoms and molecules, the study of large N-particle systems using mathematical methods, the study of quantum phases and phase transitions in strongly correlated many-body atomic and molecular systems, collisions involving highly excited states of atoms and molecules, low-energy elastic and threshold inelastic scattering of charged particles, orientation and alignment effects, the determination of potential energy surfaces, the study of atoms in magnetic and optical fields, the precision measurement of fundamental constants, the creation of more accurate and precise atomic clocks, and the generation of quantum states of light. These topics touch on many of the most important problems in physics today, such as quantum computing and quantum simulations, tests of the standard model, quantum control, and the pursuit of ever better time and frequency standards.

A large fraction of our experimental work is carried out in laboratories in Nielsen Hall. Students who choose to concentrate on experimental research work on cutting edge experiments with state-of-the-art equipment, including all forms of electronics, vacuum systems, and lasers; fiber, non-linear, electro, and acousto-optics; frequency and temperature stabilization and control. Narrow line width diode lasers are used for cooling and trapping; multiple pulsed dye lasers and pseudo continuous ultraviolet laser radiation are used in experiments designed to create ultra-cold molecules. A variety of advanced laser systems and vacuum technologies are used to investigate gases of ultracold Rydberg atoms and create entangled states of material particles. Small wires on a

chip, referred to as an atom chip, are used to trap atoms so that their interactions with a surface can be studied and utilized to construct hybrid quantum systems of atoms and surface excitations. Vapor cells and coherent multi-photon spectroscopy are used to detect small electric fields by using collections of atoms as quantum sensors. A


solid-state picosecond laser is combined with frequency stabilized diode laser radiation and microwave radiation to create an ultra-sensitive probe of molecular electric and magnetic dipole moments. Finally, narrow line width solid state lasers and atomic vapor cells are


used to generate entangled states of light for applications in quantum information and quantum metrology. This broad spectrum of experimental opportunities provides students with the training necessary to pursue careers in academia, government labs and service, and industry.

Students choosing to focus on theoretical research have many options including formal mathematical topics, very accurate and well designed physical models and large scale computational physics. Our theoretical research is at the forefront and covers a broad spectrum of current research areas (atomic physics, molecular physics, statistical physics, laser-matter interactions, chemical physics, and many-body strongly correlated systems). Our computational facilities include an extensive network of powerful computer workstations, which are freely shared among members of the department. When additional computational resources are needed the OU supercomputer was recently ranked as the 14th most powerful supercomputer at a U.S. academic institution. These facilities are used

in several experimental contexts and in theoretical research such as ongoing study of processes fundamental ultracold collision physics, coherent-control, Rydberg molecules, and large-scale Quantum Monte Carlo simulations of strongly correlated systems.

Since its inception, our group has been regularly funded by such sources as the National Science Foundation, the Department of Energy, the American Chemical Society, and the Department of Defense. In addition, various members of the group participate in long-term collaborations with scientists from Italy, France, Australia, Switzerland, Canada, Brazil, Germany, Israel, Latvia, Russia, the United Kingdom, and various laboratories and universities in the United States. A highlight of the program is a regular, intensive agenda of visits and colloquia by outside members of the atomic, molecular and chemical physics community, including our many collaborators.





[ **Eric Abraham** ]  
*associate professor*

**L. J. Semrod Presidential Professor**  
**B.A. 1991 St. Olaf College**  
**Ph.D. 1996 Rice University**

**T**he goal of our research program is to investigate ultracold atoms and molecules, including Bose-Einstein condensation. Laser cooling and trapping (the subject of the 1997 Nobel Prize) uses a variety of lasers, in addition to magnetic and electric fields, to cool atoms and molecules to a range of temperatures colder than anything else in the known universe (between 10 nanoKelvin and 100 microKelvin.) At these temperatures, their wave-like nature is enhanced allowing studies of the exotic, quantum-mechanical nature of matter.

Over 80 years ago, Albert Einstein predicted that a gas of non-interacting particles could undergo a phase transition, collecting a macroscopic number of particles into the same quantum state. The gas must be cooled to where the de Broglie wavelengths of the individual atoms overlap. This concept of Bose-Einstein condensation has since been an integral part of the understanding of strongly interacting systems such as superfluids and superconductors. However, BEC in dilute atomic gasses more accurately approximates Einstein's original prediction for non-interacting particles and its discovery was awarded the Nobel Prize in 2001. In collaboration with Deborah Watson and Michael Morrison, we are studying novel collisions and probing the strong interaction regime where connections between the gas and condensed matter systems may be found.

While laser cooling and trapping techniques have produced a revolution in atomic physics, it is limited to a few atoms. We are currently working to extend ultracold trapping techniques to molecules. In collaboration with the Shafer-Ray group at OU, we built a new apparatus that uses electric fields to produce cold gases of nitric oxide (NO). Our method uses electric fields to extract the cold fraction of particles already present in the Maxwell-Boltzmann distribution of a thermal gas. We have recently produced samples of NO in the lowest ro-vibrational quantum state at temperatures of around 1 K. This promises to open new frontiers in collision physics and ultracold chemistry. It may also lead to new systems to create quantum computers and perform precision measurements to explore physics beyond the Standard Model.

***A graph of ion signal vs laser frequency shows nitric oxide (NO) in the lowest ro-vibrational quantum state at a temperature of ~1 K. The cold gas of molecules was extracted from a thermal source of NO using static electric fields.***


T.G. Akin, S.A. Kennedy, B. Dribus, J.L. Marzoula, L. Johnson, J. Alexander, and E.R.I. Abraham, "Bose-Einstein condensation transition studies for atoms confined in Laguerre-Gaussian laser modes," *Opt. Comm.* **84**, 285 (2012).

B.J. Bichsel, M.A. Morrison, N.E. Shafer-Ray, and E.R.I. Abraham, "Experimental and theoretical investigation of the Stark effect for trapping cold molecules: Application to nitric oxide," *Phys. Rev. A* **75** 023410 (2007).

S.A. Meek, E.R.I. Abraham, and N.E. Shafer-Ray, "Impossibility of a biased Stark trap in two dimensions," *Phys. Rev. A* **71**, 065402 (2005).

E. Nikitin, E. Dashevskaya, J. Alnis, M. Auzinsh, E.R.I. Abraham, B.R. Furneaux, M. Keil, C. McRaven, N.E. Shafer-Ray, and R. Waskowsky, "Measurement and prediction of the speed-dependent throughput of a magnetic octupole velocity filter including nonadiabatic effects," *Phys. Rev. A* **68**, 023403 (2003).

S.A. Kennedy, M.J. Szabo, H. Teslow, J.Z. Porterfield, and E.R.I. Abraham, "Creation of Laguerre-Gaussian laser modes using diffractive optics," *Physical Review A* **66**, 043801 (2002).







B.S. 2000 University of Torino, Italy  
Ph.D. 2008 University of Massachusetts -  
Amherst

**M**y research is focused on many body states of matter in ultracold atomic and molecular lattice systems. Lattice systems can be realized by loading atoms or molecules into an optical lattice, i.e. a periodic array of potential wells. Such systems are highly controllable and tunable experimentally, therefore providing an excellent opportunity to test theory against experiments. At cold enough temperatures, various exotic phases can be realized in more complex systems, e.g. multi-component or dipolar ones. Often times such phases require experimentally challenging low temperatures and strong interactions. Accurate theories, capable to predict which phases and under which conditions they can be stabilized, are therefore needed.

My expertise lies in numerical methods, in particular Quantum Monte Carlo techniques. I am interested in developing new algorithms capable to efficiently treat many body systems at equilibrium, and study their properties across quantum phase transitions. I have been using the Worm Algorithm and have adapted it to multicomponent systems both, with and without, off-site interactions. The efficiency of the algorithm allows for treatment of large scale systems, and offers the opportunity for one-to-one comparison with experimental data.

Optical lattice systems have turned out to be an almost ideal experimental realization of models used to study complicated condensed matter phenomena, such as quantum magnetism or high temperature superconductivity. I am particularly interested in studying multi-component systems, relevant to quantum magnetism, with and without the presence of the external confinement potential, and dipolar systems, realized by e.g. ultracold polar molecules in the presence of an external electric field.



M. Guglielmino, V. Penna, B. Capogrosso-Sansone, "Ising antiferromagnet with ultracold bosonic mixtures confined in a harmonic trap," *Phys. Rev. A* **84**, 031603(R) (2011).

B. Capogrosso-Sansone, C. Trefzger, M. Lewenstein, P. Zoller, G. Pupillo, "Quantum Phases of Cold Polar Molecules in 2D Optical Lattices," *Phys. Rev. Lett.* **104**, 125301 (2010).

S. G. Söyler, B. Capogrosso-Sansone, N. V. Prokof'ev, and B. V. Svistunov, "Sign-Alternating Interaction Mediated by Strongly-Correlated Lattice Bosons," *New J. Phys.* **11**, 073036 (2009).

B. Capogrosso-Sansone, N. V. Prokof'ev, and B. V. Svistunov, "Phase diagram and thermodynamics of the three-dimensional Bose-Hubbard model," *Phys. Rev. B* **75**, 134302 (2007).

**Sketch of a two-dimensional optical lattice. At zero temperature, integer filling factor, i.e. integer average number of particles per site, and for deep lattices (bottom sketch), the system is a Mott insulator, that is, particles are localized and density fluctuations are suppressed. Upon decreasing the depth of potential wells, the system undergoes a quantum phase transition onto a superfluid state (upper sketch), that is, particles are delocalized, density fluctuations are enhanced, and there exist off-diagonal long range order.**



[ Alberto Marino ]  
assistant professor

B.S. 1998 ITESM, Mexico  
Ph.D. 2006 University of Rochester

**M**y main research interests lie in the fields of quantum optics and atomic physics, with particular emphasis on their applications to quantum information science and quantum metrology. Quantum optics studies the quantum properties of light and ways in which these properties can be used to provide secure communications, perform complex calculations, and increase the sensitivity of measurements, among other things.

In recent years my research has focused on the generation and control of quantum states of light known as twin beams through the use of four-wave mixing in atomic vapors. The use of an atomic system for the generation of these quantum states offers distinct advantages. First, it makes it possible to obtain a large nonlinear interaction, which leads to a large degree of entanglement without the need of a cavity. This makes the twin beams intrinsically multi-spatial mode, which means that they contain spatial quantum correlations in addition to temporal ones. Second, the use of an atomic system leads to narrowband entangled photons close to an atomic resonance. As a result, it is possible to obtain an efficient interaction between an atomic system and the entangled photons, which allows for better control of the twin beams. This has made it possible for us to implement a quantum buffer and a quantum cloner for these entangled states of light.

N.V. Corzo-Trejo, A.M. Marino, and P.D. Lett, "Noiseless Optical Amplifier Operating on Hundreds of Spatial Modes," *Phys. Rev. Lett.* **109**, 043602 (2012).

R.C. Pooser, A.M. Marino, V. Boyer, K.M. Jones, and P.D. Lett, "Low Noise Amplification of Continuous Variable Quantum States," *Phys. Rev. Lett.* **103**, 010501 (2009).


A.M. Marino, R.C. Pooser, V. Boyer, and P.D. Lett, "Tunable Delay of Einstein-Podolsky-Rosen Entanglement," *Nature* **457**, 859 (2009).

C.F. McCormick, A.M. Marino, V. Boyer, and P.D. Lett, "Strong Low-Frequency Quantum Correlations from a Four-Wave Mixing Amplifier," *Phys. Rev. A* **78**, 043816 (2008).

V. Boyer, A.M. Marino, R.C. Pooser, and P.D. Lett, "Entangled Images from Four-Wave Mixing," *Science* **321**, 544 (2008).

A.M. Marino, V. Boyer, R.C. Pooser, P.D. Lett, K. Lemons, and K.M. Jones, "Delocalized Correlations in Twin Light Beams with Orbital Angular Momentum," *Phys. Rev. Lett.* **101**, 093602 (2008).

**Quantum buffer for entangled images.** A cell containing rubidium vapor is used to generate entangled quantum images with an " $\hbar$ " shape. The dotted loop linking the images represents the entanglement between them. One of the images is sent to a second rubidium cell, which controls the propagation velocity of the light and acts as a quantum buffer. The delay can be controlled such that during the time it takes the image that goes through the quantum buffer to travel a centimeter the other image can travel up to 8 meters. Entanglement is preserved during this process.




**T**he primary emphasis of my theoretical research program in atomic, molecular, optical, and chemical physics is on purely quantum phenomena involving collisions at very low energies.

Almost all of my projects entail close collaboration with experimental physicists, both here at the University of Oklahoma and at such other institutions as the Joint Institute for Laboratory Astrophysics and the Australian National University. Typically, our research blends formal mathematics and quantum mechanics, numerical algorithms and their computational implementation, and analysis of experimental data.

Our most recent research programs include

- development of new treatment of electron correlation in scattering problems, a merger of density functional theory and many-body theory;
- a new quantum collision theory that adapts R-matrix theory (of nuclear and atomic physics) to electron transport in mesoscopic nanoscale devices;
- an alternative to conventional Boltzmann kinetic theory of electron transport in molecular gases.

At present, we are exploring the schemes for the creation of samples of ultracold molecules via processes based on photoassociation of atom-atom collisions at extremely low relative kinetic energies. Beyond this horizon, we plan to extend this research to incorporate mechanisms for quantum control that will maximize the final population of translationally and vibrationally cold molecules. This research, a collaboration with two experimentalists and a theorist in our Department, combines our long-standing interest in molecular physics and in near-threshold collision theory in contexts of great current interest to the atomic, molecular, optical, and chemical physics community.



**Photoassociation of colliding ultracold Na and Cs atoms followed by stabilization via spontaneous emission into the absolute ground state of NaCs.**


[ **Michael A. Morrison** ]  
*emeritus professor*

**American Physical Society Fellow**

**David Ross Boyd Professor**

**B.S. 1971 Rice University**

**Ph.D. 1976 Rice University**



Michael A. Morrison, **The Joy of Quantum Physics**, (Oxford University Press, 2013).

M.A. Morrison, R.E. Robson, R.D. White, **Low-energy Charged Particles in Atomic and Molecular Gases** (Princeton University Press, 2012).

Michael A. Morrison, **Understanding Quantum Physics: A User's Manual**, (Prentice-Hall 1990).

H. Feng, W. Sun, M.A. Morrison, and A.N. Feldt, "Exact inclusion of exchange in calculations of cross sections for vibrational excitation of  $N_2$  by low-energy electrons," *J. Phys. B*, **42**, 175201 (2009).

A.N. Feldt and M.A. Morrison, "The ABCs of DCS: Analytic Born completion in the calculation of differential cross sections for electron scattering from a linear molecule," *Phys. Rev. A*, **77**, 012726 (2008).

B.J. Bichsel, M.A. Morrison, N. Shafer-Ray and E.R.I. Abraham, "Experimental and theoretical investigation of the Stark effect for trapping cold molecules: application to nitric oxide," *Phys. Rev. A* **75**, 023410 (2007).

T. Jayasekera, M.A. Morrison and K. Mullen, "Cooling Electrons in Semiconductor Devices," *Phys. Rev. B* **75**, 035316 (2007).

M.A. Morrison and A.N. Feldt, "Through Scattering Theory with Gun and Camera: Coping with Conventions in Collision Theory," *Am. J. Phys.* **75**, 67 (2007).





[ Gregory A. Parker ]

*professor*

Chair of Physics and Astronomy

George Lynn Cross Research Professor

B.A. 1973 Brigham Young University

Ph.D. 1976 Brigham Young University

Over the years we have developed theoretical and numerical methods for accurately studying three-particle systems of real physical interest. Our reactive scattering theory, hyperspherical coordinates, and discrete variable representation has allowed us and others to solve the Schrodinger equation with minimal approximations. In addition we have also developed rotational decoupling approximations which are widely used in atom-diatom inelastic scattering and has led to ultrasimple expressions for calculating integral and differential cross sections.

We have studied a variety of systems including: Positronium formation which occurs when a positron collides with a Hydrogen atom.

Positronium is an exotic atom composed of a positron-electron pair. The highly exoergic  $F+H_2$  reaction is an important bottle neck step in powerful hydrogen fluoride lasers. We also studied the single most important combustion reaction  $H+O_2$ . This reaction is the rate limiting step which determines rates of explosion and flame propagation.

Currently, we are interested in ultracold collisions of alkali atoms with alkali dimers. This interest is the result of the phenomenal success in the experimental formation of ultracold atoms and molecules. We have recently shown that pulsed lasers of moderate intensities used during the collision can lead to the efficient production of ultracold molecules. This laser catalyzed production of ultracold molecules utilizes a laser to quantum mechanically control the chemical reaction. This is accomplished by forcing a virtual transition of the reactants to an excited state complex. Then the excited state complex undergoes stimulated emission back to the ground electronic state, releasing a photon identical to the absorbed photon.

Another reason for the interest in triatomic  $Li_3$  is the existence of energetically accessible conical intersections in both the doublet and quartet states. A conical intersection is seen where two different Born-Oppenheimer electronic states intersect. The presence of conical intersections affect the bound states of  $Li_3$  and the  $Li+Li_2$  dynamics.

We are also interested in collisions in the presence of intense laser fields. Reactive collisions occur at relatively short internuclear distances which strongly couples with the intense laser field drastically altering the dynamics.

X. Li, W. Dupre, G.A. Parker, "Pulse Sequences in Photoassociation via Adiabatic Passage," *New J. Phys.* **14**, 073001 (2012).


X. Li, D.A. Brue, B.K. Kendrick, J.D. Bandon, G.A. Parker, "Geometric Phase for Collinear Conical Intersections. I. Geometric Phase Angle and Vector Potentials. *J. Chem. Phys.* **134**, 064108 (2011).

X. Li, G.A. Parker, P. Brumer, I. Thanopoulos, M. Shapiro, "Laser-Catalyzed Production of Ultracold Molecules: The  ${}^6Li + {}^6Li \xrightarrow{h\nu} {}^6Li_2 + {}^7Li$  Reaction." *Phys. Rev. Lett.* **101**, 043003 (2008).

X. Li, D.A. Brue, G.A. Parker, "Potential Energy Surfaces for  $1^4A'$ ,  $2^4A'$ ,  $1^4A''$ ,  $2^4A''$  States of  $Li_3$ . Use of Diatomics in Molecules to Fit Full Configuration Interaction Data." *J. Chem. Phys.* **129**, 124305 (2008).

X. Li, G.A. Parker, "Theory of Laser Enhancement of Ultracold Reactions: The Fermion-Boson Population Transfer Adiabatic Passage of  ${}^6Li + {}^6Li \xrightarrow{h\nu} {}^6Li_2 + {}^7Li$  ( $T_f=1mK$ )  $\rightarrow {}^6Li_2 + {}^7Li$  ( $T_f=1mK$ )," *J. Chem. Phys.* **128**, 184113 (2008).


X. Li, G.A. Parker, P. Brumer, I. Thanopoulos, M. Shapiro, "Theory of Laser Enhancement and Suppression of Cold Reactions: The Fermion-Boson  ${}^6Li+{}^7Li_2 \xleftrightarrow{h\nu} {}^6Li \xrightarrow{h\nu} {}^6Li+{}^7Li$  Radiative Collision." *J. Chem. Phys.* **128**, 124314 (2008).



**Three-dimensional view of the potential energy surface in the  $C_{2v}$  plane. This conical intersection is the lowest point on the seam of conical for the spin-aligned state of  $Li_3$ .**

[ Neil Shafer-Ray ]  
professor

B.S. 1986 M.I.T.  
Ph.D. 1990 Columbia University



P. Sivakumar, C.P. McRaven, T. Yang, M. Rupasinghe, N.E. Shafer-Ray, T. Sears, and G. Hall, "Pseudo-continuous resonance enhanced multiphoton ionization: Application to the determination of the hyperfine constants of  $^{208}\text{Pb}^{19}\text{F}$ ," *Journal of Molecular Dynamics In Preparation* (2009).

C.P. McRaven, P. Sivakumar, and N.E. Shafer-Ray, "Experimental determination of the hyperfine states of the  $^{207}\text{Pb}^{19}\text{F}$  molecule," *Phys. Rev. A*, 054502 (2008).

M. Rupasinghe and N.E. Shafer-Ray, "Effect of the geometric phase on the possible measurement of the electron's electric dipole moment using molecules confined by a Stark gravitational trap," *Phys. Rev. A* **78**, 033427 (2008).

Bryan J. Bichsel, Michael A. Morrison, Neil Shafer-Ray, and E. R. Abraham, "Experimental and theoretical investigation of the Stark effect for manipulating cold molecules: Application to nitric oxide," *Phys. Rev. A* **75**, 023410 (2007).

Neil Shafer-Ray, "Possibility of zero-g-factor paramagnetic molecules for measurement of the electron's electric dipole moment," *Phys. Rev. A* **73**, 034102 (2006).


Our group currently covers three research areas: The first is an effort to use a novel resonance technique to measure the electron's electric dipole moment. The second is an effort to use recently developed techniques in cooling and trapping to create a source of ultracold nitric oxide molecules.

It is well known that an electron carries a magnetic moment  $\vec{m} = \mu_B g \vec{S}$  that is proportional to its spin. (Here  $\mu_B$  is the Bohr magneton and  $g$  the dimensionless "g-factor".) In the early 1950's Purcell and Ramsey suggested that the electron may also carry an electric dipole moment leading to an interaction

$$U = -\mu_B (g \vec{S} \cdot \vec{B} + g_{EDM} \vec{S} \cdot \vec{E} / c)$$

Whereas the  $g$  factor is known to be a bit larger than 2, current experiments limit  $|g_{EDM} \mu_B / c|$  to less than  $10^{-27}$  e cm. A major focus of our group is to either measure or place an even smaller limit on the value of  $g_{EDM}$ . To do so we are observing magic values of electric field for which the magnet moment of the molecule  $\text{PbF}$  vanishes. If the electron has a dipole moment, this electric field value will be different for the case of a magnetic field parallel to an electric field and the case of a magnetic field anti-parallel to an electric field. Measurement of a non-zero electric dipole moment would help differentiate between competing models of Particle Physics and could help explain mechanisms for CP violation that might have led to a matter-dominated Universe.

Cold populations of gas phase molecules contained by non-uniform electro-magnetic fields are of great importance to many fields, including precision measurement, navigation, time standards, and quantum computing. Our group is working with the Abraham laboratory to create an ultracold source of nitric oxide molecules. Here slow moving nitric oxide molecules are optically pumped into a trap state using pseudo continuous laser radiation. This creates a relatively hot ( $\sim 1$  K) population of trapped molecules. The temperature is then reduced with a second optical pumping scheme (demonstrated for atoms by the Raizen group at the University of Texas Austin.) The goal of this research is to create a dense source of NO molecules at a temperature below 1 mK.



**A electric dipole moment  $p$  of the electron would give a length scale  $p/e$  to the electron. The figure illustrates the relative size of the current limit of this scale ( $10^{-14}$  fm). The goal of our electron electric dipole moment project is to probe for even smaller values of  $p/e$ .**



[ James Shaffer ]  
*associate professor*

**B.S. 1991 University of Illinois  
 at Urbana-Champaign**

**Ph.D. 1999 University of Rochester**

“Atom Optics: Marriage of Atoms and Plasmons,” J.P. Shaffer, *Nature Photonics* **5**, 451 (2011).

“Coherent Excitation of Rydberg Atoms in Thermal Vapor Cells,” H. Kubler, J.P. Shaffer, T. Baluksian, R. Low, and T. Pfau, *Nature Photonics* **4**, 112 (2010).

“Electric Field Effects in the Excitation of Cold Rydberg Atom Pairs,” V.A. Nascimento, L.L. Caliri, A. Schwettmann, J.P. Shaffer, and L.G. Marcassa, *Phys. Rev. Lett.* **102**, 213201 (2009).

“Observation of Ultra-Long Range Rydberg Molecules,” V. Bendkowsky, B. Butscher, J. Nipper, J.P. Shaffer, R. Low, and T. Pfau, *Nature* **458**, 1005 (2009).

“Observation of Electric-Field Induced Cs Rydberg Atom Macrodimers,” K.R. Overstreet, A. Schwettmann, J. Tallant, D. Booth, and J.P. Shaffer, *Nature Physics* **5**, 581-585 (2009).

I am currently interested in investigating exotic states of matter and the coupling of quantum dynamical systems to their environment. My research group has developed novel experimental methods for investigating the interactions between ultracold atoms using Rydberg atom time-of-flight spectroscopy. In developing this method, we were able to predict and find a new type of molecule formed by 2 Rydberg atoms which has a bond length of  $> 3 \mu\text{m}$ . I am also interested in other types of molecules that can form at ultralow temperatures such as Rydberg atom-ground state atom molecules and the physics of Efimov states and three-body recombination. These types of systems and methods are important for investigating chemical dynamics that can take place in ultracold atom traps. My research group also is working on understanding the interactions between cold Rydberg atoms for making new types of quantum entanglement based devices such as single photon sources and quantum gates. Some of this work is done at ultracold temperatures ( $T \leq 1 \text{ mK}$ ) and some is done in micron-sized thermal vapor cells. Most of these investigations are experimental but my research group also does theory to support our experimental efforts.



*Rydberg atom imaging spectrometer.*



[ Deborah K. Watson ]  
*professor*

Edith Kinney Gaylord Presidential Professor  
B.A. 1972 Allegheny College  
Ph.D. 1977 Harvard

**M**y group is engaged in the study of large N-particle systems under quantum confinement such as Bose-Einstein condensates. An exact solution of the N-body problem is considered to be an “intractable” problem, scaling exponentially with the number of particles, N. The resources required for a solution typically double with every particle added, resulting in a current limit of ~10 particles for an exact solution. We are developing a method which circumvents this exponential scaling with N by rearranging the work so the problem now scales exponentially with order in a perturbation series. Exact solutions are possible at each order for any N. We have accomplished this by using group theoretic as well as graphical techniques. These powerful and elegant techniques “hold their own” as N increases resulting in minimum numerical effort.

We are presently pursuing several studies including the calculation of both ground and excited state energies for Bose-Einstein condensates using trap parameters that approximate current experimental conditions, the determination of an analytic many-body density profile from the first-order wavefunction and an analysis of the fundamental motions associated with excitation. A Bose-Einstein condensate is an ideal system to test a new many-body approach since it is a coherent macroscopic sample of atoms in the same quantum state.

In addition, we have started the extension of our formalism to large systems of strongly interacting fermions which are important in many fields of physics including astrophysics, nuclear, condensed matter as well as atomic physics. Our goal is to bridge the gap between a microscopic quantum Hamiltonian and the macroscopic properties of large N-particle correlated systems.



D.K. Watson and M. Dunn, “Analysis of the growth in complexity of a symmetry-invariant perturbation method for large N-body systems,” *J. Phys. B* **45**, 095002 (2012).

D.K. Watson and M. Dunn, “Rearranging the Exponential Wall for Large N-Body Systems,” *Phys. Rev. Lett.* **105**, 020402 (2010).

M. Dunn, W. B. Laing, D. Toth, and D.K. Watson, “A Test of a New Interacting N-Body Wave Function,” *Phys. Rev. A* **80**, 062108 (2009).

W.B. Laing, D.W. Kelle, M. Dunn, and D.K. Watson, “A Complete Basis for a Perturbation of the General N-Body Problem,” *J. Phys. A* **42**, 205307 (2009).

W.B. Laing, M. Dunn, and D.K. Watson, “On the Use of Group Theoretical and Graphical Techniques to solve the N-Body Problem with General Two-Body Interactions,” *J. Math. Phys.* **50**, 062105 (2009).

W.B. Laing, M. Dunn, and D.K. Watson, “Analytic, group-theoretic density profiles for confined, correlated N-body systems,” *Phys. Rev. A* **74**, 06360 (2006).

M. Dunn, D.K. Watson and J.G. Loeser, “Analytic, group theoretic wave functions for confined, correlated N-body systems with general two-body interactions,” *Ann. Phys.* **321**, 1939 (2006).